

## What it does

When soil experience crusting, the germinating seeds are unable to break the crust. This also limits oxygen flow into and out of the soil, thus reducing crop growth.

## Why and where does it occur

Crusting occurs as the soil dries.

The problem is primarily in dry direct-seeded fields where seed is covered by soil, but may also occur in wet direct-seeded fields if the soil dries during crop emergence.

## How to identify

Check the field for the following symptoms:

- Poor crop emergence in direct seeded fields
- Plants fail to emerge, as they cannot break through the drying soil surface

The pattern of damage depends on the leveling of the field and the pattern of soil drying across the field.

Various factors that may cause problems of crusting are soil type, high intensity heavy rainfall at seeding, and dry weather subsequent to rains.

To confirm cause of problem, check or ask farmer if soil has sealed (become hard) at the time of crop emergence in direct-seeded field. Also check for plants that have germinated but roots have not been able to penetrate the soil surface so well.

## Why is it important

Good planting or crop establishment lays the foundation for good yield. Crusting greatly reduces crop stand in direct seeded fields. Its economic effect is direct due to a reduced plant stand and subsequent yield reduction.

## How to manage

Crusting is primarily important just at the time of crop emergence.

If the soil surface can be kept moist, then crusting is unlikely to be a problem.

- Keep the soil surface moist.
- Maintain residue at soil surface. It would decelerate the falling raindrops, and thereby reduce their impact on soil; also it would prevent evaporation of moisture from top soil.
- Apply irrigation well before crusting starts or it has just initiated.

